

# *Regional Results*

## *2001*

### *Region I*

*Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, Vermont*

### *Region II*

*New Jersey, New York*

### *Region III*

*Delaware, District of Columbia, Maryland, Pennsylvania,  
Virginia, West Virginia*

### *Region IV*

*Alabama, Florida, Georgia, Kentucky, Mississippi, North Carolina,  
South Carolina, Tennessee*

### *Region V*

*Illinois, Indiana, Michigan, Minnesota, Ohio, Wisconsin*

### *Region VI*

*Arkansas, Louisiana, New Mexico, Oklahoma, Texas*

### *Region VII*

*Iowa, Kansas, Missouri, Nebraska*

### *Region VIII*

*Colorado, Montana, North Dakota, South Dakota, Utah, Wyoming*

### *Region IX*



*Arizona, California, Hawaii, Nevada*

### *Region X*

*Alaska, Idaho, Oregon, Washington*

# Region I – CT, ME, MA, NH, RI, VT



Resident Characteristics*	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
Nursing Home Residents	109,657	90,192	6,758	7,092	4,915
Percent of Nursing Home Residents	100.00	82.25	6.16	6.47	4.48
Percent Female	74.69	74.85	75.41	72.70	73.87
Age: <25	0.28	0.25	0.59	0.11	0.59
Age: 25-54	3.13	3.26	2.68	2.26	2.77
Age: 55-64	3.55	3.65	2.93	3.44	2.85
Age: 65-74	9.59	9.53	10.83	9.74	8.73
Age: 75-84	30.56	30.58	29.55	31.91	29.77
Age: 85-94	42.44	42.38	42.99	41.51	43.76
Age: 95+	10.45	10.35	10.43	11.03	11.54
American Indian/Alaskan Native	0.20	0.19	0.33	0.20	0.20
Asian/Pacific Islander	0.43	0.49	0.13	0.13	0.10
Black	3.55	4.17	0.52	0.87	0.37
Hispanic	1.30	1.53	0.41	0.18	0.12
White	94.52	93.61	98.61	98.62	99.21
Lived Alone	31.96	32.40	28.33	32.53	26.59
Lived with Others	51.07	50.63	55.48	53.42	50.70
Lived in Other Facility	16.96	16.97	16.19	14.05	22.71
Admitted from Private Home with No Services	11.63	11.70	12.31	10.50	11.52
Admitted from Private Home with Services	8.82	8.38	10.55	11.20	12.76
Admitted from Assisted Living	4.79	4.51	5.75	6.36	7.48
Admitted from Other Nursing Home	15.22	15.57	14.55	13.00	12.42
Admitted from Acute Care Hospital	51.92	52.14	50.45	53.67	45.31
Admitted from Other Setting	7.61	7.70	6.40	5.27	10.51
CPS: Intact or Borderline Intact	22.46	22.85	22.28	20.76	18.72
CPS: Mild to Moderately Severely Impaired	52.73	52.54	51.07	54.71	54.97
CPS: Severely or Very Severely Impaired	24.81	24.61	26.65	24.53	26.31
ADL: Independent or Needs Supervision	11.31	11.96	9.87	7.26	7.37
ADL: Limited Physical Assistance	13.24	13.55	13.76	10.86	10.05
ADL: Extensive Assistance in Early Loss ADLs	22.58	22.56	22.77	22.01	23.87
ADL: Extensive Assistance in Later Loss ADLs	12.10	11.16	14.86	18.27	17.05
ADL: Totally Dependent in Some ADLs	24.45	24.06	24.40	27.19	27.08
ADL: Totally Dependent in All Four ADLs	16.32	16.71	14.34	14.41	14.59



## Region I 2001 (continued)



	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
CHES: No Instability	35.89	37.29	29.61	28.47	28.95
CHES: Slightly Unstable	30.55	30.24	32.73	31.94	31.03
CHES: Mildly Unstable	18.52	17.86	21.38	22.03	21.71
CHES: Moderately Unstable	10.11	9.85	10.85	11.79	11.72
CHES: Highly Unstable	4.93	4.76	5.43	5.77	6.59
Total Case-Mix Index (average)	0.87	0.87	0.88	0.89	0.91
Nursing Case-Mix Index (average)	0.81	0.81	0.81	0.84	0.83
<b>Quality Indicators</b>					
Depression without Therapy	6.12	5.48	7.61	9.25	11.77
9+ Different Medications	47.23	46.96	48.51	49.27	48.40
Bladder/Bowel Incontinence without Plan	34.00	33.17	40.01	36.57	32.01
Indwelling Catheters	7.92	7.97	7.74	8.53	6.71
Tube Feeding	4.38	4.65	3.48	2.96	2.64
Antipsychotic Use without Conditions	21.20	21.61	19.60	18.87	18.97
Antianxiety/Hypnotic Use	18.13	17.96	20.20	18.29	19.29
Daily Physical Restraint	9.27	9.75	5.76	5.50	9.64
Falls	17.26	16.91	18.32	19.11	19.59
Behavioral Symptoms Affecting Others	21.35	20.73	22.08	23.67	28.25
Symptoms of Depression	14.11	12.99	16.43	19.68	24.46
Bladder/Bowel Incontinence	62.86	62.50	62.41	64.79	66.27
Fecal Impaction	0.11	0.10	0.18	0.18	0.06
Urinary Tract Infections	10.58	10.57	10.42	11.35	9.89
Weight Loss	15.65	15.55	15.49	16.86	16.55
Dehydration	1.29	1.24	1.88	1.52	1.08
Bedfast Residents	4.20	4.07	4.22	5.23	4.84
Little/No Activity	13.25	13.12	13.36	13.12	16.16
Stage 1-4 Pressure Ulcer	16.22	16.66	14.80	13.74	14.51

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.



## Region II – NJ, NY



Resident Characteristics*	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
Nursing Home Residents	159,502	140,289	8,683	8,103	2,427
Percent of Nursing Home Residents	100.00	87.95	5.44	5.08	1.52
Percent Female	72.00	71.78	74.88	72.02	74.25
Age: <25	0.49	0.54	0.14	0.04	0.21
Age: 25-54	4.02	4.12	4.43	2.23	2.88
Age: 55-64	4.68	4.82	3.80	3.43	3.71
Age: 65-74	11.08	11.41	8.13	9.10	9.35
Age: 75-84	30.19	30.03	31.18	32.03	29.71
Age: 85-94	39.67	39.33	41.29	43.07	42.52
Age: 95+	9.87	9.76	11.03	10.11	11.62
American Indian/Alaskan Native	0.32	0.23	0.31	2.00	0.21
Asian/Pacific Islander	1.16	1.31	0.12	0.14	0.04
Black	13.55	15.10	2.06	2.67	1.69
Hispanic	4.73	5.31	0.51	0.64	0.29
White	80.23	78.06	97.00	94.56	97.78
Lived Alone	27.21	27.34	28.74	24.65	21.94
Lived with Others	60.13	60.42	57.69	57.63	61.68
Lived in Other Facility	12.66	12.24	13.57	17.73	16.39
Admitted from Private Home with No Services	12.21	12.25	9.57	14.26	13.29
Admitted from Private Home with Services	5.15	4.88	7.61	5.75	8.77
Admitted from Assisted Living	4.73	4.38	6.18	8.01	7.74
Admitted from Other Nursing Home	10.45	10.51	10.42	8.33	14.45
Admitted from Acute Care Hospital	63.11	63.59	61.51	60.30	50.71
Admitted from Other Setting	4.36	4.38	4.71	3.35	5.03
CPS: Intact or Borderline Intact	25.06	25.06	26.65	23.14	26.33
CPS: Mild to Moderately Severely Impaired	51.06	51.11	51.09	50.50	49.90
CPS: Severely or Very Severely Impaired	23.88	23.83	22.26	26.36	23.77
ADL: Independent or Needs Supervision	11.19	11.51	9.43	8.43	8.03
ADL: Limited Physical Assistance	12.67	12.68	12.45	12.35	13.84
ADL: Extensive Assistance in Early Loss ADLs	20.61	20.50	20.53	22.26	21.80
ADL: Extensive Assistance in Later Loss ADLs	9.19	9.05	10.48	9.85	10.22
ADL: Totally Dependent in Some ADLs	24.63	24.19	28.98	27.08	26.33
ADL: Totally Dependent in All Four ADLs	21.72	22.07	18.13	20.03	19.78



## Region II 2001 (continued)



	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
CHES: No Instability	42.98	44.43	31.95	32.50	33.35
CHES: Slightly Unstable	29.58	29.10	33.68	32.49	32.81
CHES: Mildly Unstable	16.12	15.61	19.62	20.11	20.03
CHES: Moderately Unstable	7.78	7.54	9.70	9.60	8.86
CHES: Highly Unstable	3.54	3.32	5.06	5.31	4.95
Total Case-Mix Index (average)	0.89	0.89	0.85	0.87	0.90
Nursing Case-Mix Index (average)	0.82	0.82	0.82	0.82	0.85
<b>Quality Indicators</b>					
Depression without Therapy	4.14	3.95	4.85	5.86	6.46
9+ Different Medications	36.79	35.81	45.36	41.75	46.07
Bladder/Bowel Incontinence without Plan	36.96	36.36	41.94	40.00	34.33
Indwelling Catheters	7.28	7.14	8.45	7.70	9.48
Tube Feeding	10.22	10.87	5.16	5.37	6.84
Antipsychotic Use without Conditions	18.54	18.46	18.36	20.02	19.18
Antianxiety/Hypnotic Use	11.46	11.18	14.10	12.69	13.50
Daily Physical Restraint	7.86	7.53	9.55	10.49	12.44
Falls	13.57	13.20	16.30	16.73	14.79
Behavioral Symptoms Affecting Others	18.11	17.92	19.31	19.39	20.12
Symptoms of Depression	8.13	7.67	10.14	12.35	13.50
Bladder/Bowel Incontinence	63.50	63.46	62.52	66.03	60.78
Fecal Impaction	0.19	0.18	0.30	0.20	0.04
Urinary Tract Infections	8.74	8.57	9.72	10.22	10.14
Weight Loss	15.94	15.98	15.87	15.20	16.12
Dehydration	0.83	0.81	0.93	1.02	0.70
Bedfast Residents	3.67	3.64	4.25	3.76	3.58
Little/No Activity	10.43	10.21	12.50	11.60	11.76
Stage 1-4 Pressure Ulcer	23.89	24.69	17.86	18.99	19.63

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.



# Region III – DE, DC, MD, PA, VA, WV



Resident Characteristics*	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
Nursing Home Residents	181,103	131,427	15,073	21,370	13,110
Percent of Nursing Home Residents	100.00	72.57	8.32	11.80	7.24
Percent Female	73.78	73.90	74.22	73.21	72.96
Age: <25	0.15	0.18	0.04	0.11	0.09
Age: 25-54	3.44	3.66	2.95	2.75	2.92
Age: 55-64	4.11	4.17	3.79	4.11	3.87
Age: 65-74	11.19	11.16	11.31	11.28	11.15
Age: 75-84	33.20	33.30	33.17	32.91	32.74
Age: 85-94	39.25	38.82	40.47	40.36	40.32
Age: 95+	8.67	8.71	8.28	8.49	8.90
American Indian/Alaskan Native	0.11	0.11	0.11	0.15	0.08
Asian/Pacific Islander	0.43	0.52	0.25	0.14	0.11
Black	14.87	18.09	6.32	6.01	7.01
Hispanic	0.54	0.68	0.17	0.10	0.17
White	84.05	80.60	93.15	93.59	92.63
Lived Alone	24.26	23.74	25.49	26.49	23.33
Lived with Others	57.59	58.90	53.66	54.87	56.75
Lived in Other Facility	18.15	17.36	20.85	18.64	19.92
Admitted from Private Home with No Services	14.48	13.50	14.18	15.66	19.28
Admitted from Private Home with Services	4.91	4.51	4.78	5.65	6.49
Admitted from Assisted Living	7.93	7.90	8.21	7.29	8.76
Admitted from Other Nursing Home	11.54	11.07	13.30	12.26	11.90
Admitted from Acute Care Hospital	55.23	57.46	54.16	52.19	46.43
Admitted from Other Setting	5.91	5.57	5.37	6.93	7.13
CPS: Intact or Borderline Intact	22.67	22.70	22.52	23.89	20.70
CPS: Mild to Moderately Severely Impaired	50.04	50.08	50.82	48.36	51.44
CPS: Severely or Very Severely Impaired	27.29	27.22	26.66	27.75	27.86
ADL: Independent or Needs Supervision	7.27	7.32	6.36	7.46	7.52
ADL: Limited Physical Assistance	10.14	10.18	9.26	10.44	10.19
ADL: Extensive Assistance in Early Loss ADLs	21.05	21.04	22.14	20.58	20.77
ADL: Extensive Assistance in Later Loss ADLs	14.07	13.49	15.86	15.01	16.24
ADL: Totally Dependent in Some ADLs	26.63	26.61	26.82	27.08	25.96
ADL: Totally Dependent in All Four ADLs	20.84	21.37	19.56	19.42	19.31



## Region III 2001 (continued)



	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
CHES: No Instability	31.41	32.88	27.90	26.57	28.71
CHES: Slightly Unstable	31.18	30.93	32.69	31.46	31.46
CHES: Mildly Unstable	20.90	20.32	21.96	23.03	22.03
CHES: Moderately Unstable	11.19	10.78	11.81	12.84	11.86
CHES: Highly Unstable	5.32	5.10	5.63	6.10	5.93
Total Case-Mix Index (average)	1.00	1.00	1.01	1.03	0.98
Nursing Case-Mix Index (average)	0.90	0.89	0.90	0.91	0.89
<b>Quality Indicators</b>					
Depression without Therapy	6.99	6.63	7.10	7.88	9.09
9+ Different Medications	50.93	50.11	52.13	54.83	51.45
Bladder/Bowel Incontinence without Plan	53.52	53.95	51.83	54.63	50.41
Indwelling Catheters	11.09	11.12	10.36	11.45	11.17
Tube Feeding	9.94	10.98	7.56	6.91	7.19
Antipsychotic Use without Conditions	17.68	17.70	17.66	17.79	17.34
Antianxiety/Hypnotic Use	17.24	16.76	18.54	18.88	17.94
Daily Physical Restraint	7.75	7.72	8.66	7.26	7.80
Falls	17.63	17.36	18.39	17.99	18.83
Behavioral Symptoms Affecting Others	20.12	19.18	20.53	22.39	25.51
Symptoms of Depression	14.45	13.88	14.76	15.74	17.69
Bladder/Bowel Incontinence	68.75	69.43	68.33	66.34	66.37
Fecal Impaction	0.27	0.24	0.28	0.30	0.41
Urinary Tract Infections	13.27	13.27	12.67	13.67	13.30
Weight Loss	17.83	18.04	17.16	18.00	16.04
Dehydration	1.14	1.12	1.12	1.14	1.38
Bedfast Residents	7.53	7.54	7.45	7.84	6.96
Little/No Activity	15.41	15.45	14.36	15.83	15.51
Stage 1-4 Pressure Ulcer	21.13	22.32	17.95	18.32	18.03

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.



# Region IV– AL, FL, GA, KY, MS, NC, SC, TN



Resident Characteristics*	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
Nursing Home Residents	292,619	171,694	44,775	51,818	23,116
Percent of Nursing Home Residents	100.00	58.67	15.30	17.71	7.90
Percent Female	73.47	73.12	74.49	73.90	73.14
Age: <25	0.24	0.29	0.19	0.14	0.16
Age: 25-54	3.50	3.72	2.93	3.33	3.41
Age: 55-64	4.74	4.73	4.58	4.70	5.24
Age: 65-74	12.33	12.30	12.51	12.54	11.91
Age: 75-84	32.81	32.62	33.36	33.10	32.53
Age: 85-94	38.19	37.96	38.52	38.59	38.28
Age: 95+	8.19	8.38	7.92	7.61	8.48
American Indian/Alaskan Native	0.19	0.13	0.25	0.21	0.45
Asian/Pacific Islander	0.17	0.23	0.08	0.07	0.04
Black	19.30	20.10	20.26	17.20	16.77
Hispanic	2.01	3.24	0.26	0.26	0.16
White	78.33	76.29	79.14	82.25	82.57
Lived Alone	21.37	20.06	24.49	23.17	20.86
Lived with Others	64.08	64.36	62.20	63.85	66.52
Lived in Other Facility	14.55	15.58	13.31	12.99	12.62
Admitted from Private Home with No Services	14.44	13.25	13.57	17.24	19.24
Admitted from Private Home with Services	4.46	3.46	5.09	5.83	7.85
Admitted from Assisted Living	5.90	6.89	5.09	4.29	3.51
Admitted from Other Nursing Home	12.62	12.44	11.01	13.93	14.38
Admitted from Acute Care Hospital	57.14	58.12	61.04	53.40	49.76
Admitted from Other Setting	5.43	5.83	4.19	5.31	5.25
CPS: Intact or Borderline Intact	20.61	20.33	21.44	20.92	20.67
CPS: Mild to Moderately Severely Impaired	52.36	52.94	52.36	50.79	51.28
CPS: Severely or Very Severely Impaired	27.03	26.72	26.20	28.29	28.04
ADL: Independent or Needs Supervision	10.25	10.43	10.28	10.20	9.05
ADL: Limited Physical Assistance	13.01	13.06	13.08	12.53	13.54
ADL: Extensive Assistance in Early Loss ADLs	19.98	20.55	19.07	19.04	19.45
ADL: Extensive Assistance in Later Loss ADLs	10.22	10.09	10.39	10.12	10.95
ADL: Totally Dependent in Some ADLs	23.38	23.51	23.59	23.22	22.43
ADL: Totally Dependent in All Four ADLs	23.16	22.36	23.59	24.89	24.58



## Region IV 2001 (continued)



	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
CHES: No Instability	31.31	31.47	32.01	30.99	29.64
CHES: Slightly Unstable	30.63	30.31	30.87	31.27	31.26
CHES: Mildly Unstable	21.42	21.31	21.30	21.54	22.25
CHES: Moderately Unstable	11.35	11.46	10.84	11.21	11.61
CHES: Highly Unstable	5.29	5.44	4.98	4.99	5.23
Total Case-Mix Index (average)	0.97	0.98	0.95	0.94	0.93
Nursing Case-Mix Index (average)	0.85	0.85	0.85	0.85	0.86
<b>Quality Indicators</b>					
Depression without Therapy	5.70	5.37	5.83	6.09	7.04
9+ Different Medications	48.61	47.26	49.08	51.19	52.00
Bladder/Bowel Incontinence without Plan	49.59	46.96	57.33	52.61	49.39
Indwelling Catheters	10.15	10.24	9.76	10.22	10.16
Tube Feeding	11.60	11.91	11.27	11.03	11.27
Antipsychotic Use without Conditions	18.84	18.00	19.13	20.46	20.95
Antianxiety/Hypnotic Use	20.38	19.39	20.67	22.40	22.62
Daily Physical Restraint	10.76	10.15	11.06	11.77	12.23
Falls	16.34	16.57	16.19	15.87	15.92
Behavioral Symptoms Affecting Others	19.48	18.75	19.21	20.82	22.14
Symptoms of Depression	11.87	11.40	11.64	12.52	14.12
Bladder/Bowel Incontinence	64.80	65.90	63.70	63.15	62.17
Fecal Impaction	0.39	0.39	0.47	0.38	0.33
Urinary Tract Infections	12.83	13.36	11.32	12.48	12.47
Weight Loss	20.00	20.34	19.69	19.38	19.34
Dehydration	1.14	1.12	1.18	1.16	1.17
Bedfast Residents	10.79	9.63	12.18	12.61	12.94
Little/No Activity	16.93	16.24	18.19	17.50	18.62
Stage 1-4 Pressure Ulcer	19.21	20.60	17.37	17.22	17.09

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.



# Region V – IL, IN, MI, MN, OH, WI



Resident Characteristics*	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
Nursing Home Residents	355,035	231,747	50,463	42,962	29,200
Percent of Nursing Home Residents	100.00	65.27	14.21	12.10	8.22
Percent Female	72.13	72.06	72.67	73.20	70.19
Age: <25	0.18	0.15	0.45	0.09	0.07
Age: 25-54	4.67	5.52	3.76	2.59	2.57
Age: 55-64	4.69	5.28	4.10	3.48	2.81
Age: 65-74	10.83	11.46	10.16	9.47	8.98
Age: 75-84	30.81	30.96	31.09	30.44	29.76
Age: 85-94	39.52	37.84	40.81	43.45	44.86
Age: 95+	9.30	8.80	9.63	10.48	10.95
American Indian/Alaskan Native	0.24	0.22	0.19	0.23	0.57
Asian/Pacific Islander	0.37	0.52	0.11	0.04	0.05
Black	9.30	13.48	2.10	1.10	0.65
Hispanic	0.71	0.95	0.37	0.16	0.21
White	89.38	84.83	97.22	98.47	98.52
Lived Alone	28.59	26.80	31.95	33.80	31.60
Lived with Others	53.99	54.95	51.69	50.59	54.38
Lived in Other Facility	17.42	18.25	16.36	15.61	14.03
Admitted from Private Home with No Services	16.25	14.73	17.70	20.97	20.98
Admitted from Private Home with Services	4.74	4.04	5.73	6.02	7.70
Admitted from Assisted Living	5.99	6.03	6.49	5.26	5.69
Admitted from Other Nursing Home	15.97	16.12	15.92	15.04	16.11
Admitted from Acute Care Hospital	50.88	52.59	49.19	46.33	44.47
Admitted from Other Setting	6.17	6.48	4.98	6.38	5.05
CPS: Intact or Borderline Intact	23.30	22.50	24.25	25.17	25.38
CPS: Mild to Moderately Severely Impaired	56.23	57.12	54.86	54.65	53.85
CPS: Severely or Very Severely Impaired	20.47	20.38	20.90	20.18	20.77
ADL: Independent or Needs Supervision	15.24	15.65	13.77	14.52	15.65
ADL: Limited Physical Assistance	14.51	14.58	14.31	14.82	13.96
ADL: Extensive Assistance in Early Loss ADLs	22.19	21.79	22.20	22.78	24.52
ADL: Extensive Assistance in Later Loss ADLs	13.44	12.96	14.63	14.18	13.89
ADL: Totally Dependent in Some ADLs	20.70	20.73	21.40	20.64	19.47
ADL: Totally Dependent in All Four ADLs	13.92	14.29	13.69	13.07	12.52



## Region V 2001 (continued)



	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
CHES: No Instability	30.75	32.50	25.91	27.78	29.52
CHES: Slightly Unstable	31.24	30.74	32.44	31.88	32.24
CHES: Mildly Unstable	21.32	20.68	23.67	22.25	20.98
CHES: Moderately Unstable	11.30	10.92	12.32	12.08	11.36
CHES: Highly Unstable	5.39	5.16	5.66	6.01	5.89
Total Case-Mix Index (average)	0.93	0.94	0.94	0.90	0.87
Nursing Case-Mix Index (average)	0.83	0.83	0.84	0.82	0.81
<b>Quality Indicators</b>					
Depression without Therapy	9.43	9.08	10.06	10.45	9.69
9+ Different Medications	50.78	49.95	53.51	52.05	50.59
Bladder/Bowel Incontinence without Plan	31.31	32.66	30.55	29.32	26.92
Indwelling Catheters	9.42	9.68	9.52	9.18	7.57
Tube Feeding	6.55	7.95	4.61	3.60	3.05
Antipsychotic Use without Conditions	17.72	18.32	17.35	16.67	15.15
Antianxiety/Hypnotic Use	17.50	16.79	19.73	19.34	16.31
Daily Physical Restraint	7.65	7.50	8.59	7.60	7.20
Falls	18.47	17.86	19.22	20.06	19.71
Behavioral Symptoms Affecting Others	27.06	26.68	27.20	28.16	28.30
Symptoms of Depression	19.49	19.02	20.59	20.65	19.77
Bladder/Bowel Incontinence	55.60	55.93	55.04	55.28	54.32
Fecal Impaction	0.16	0.17	0.16	0.17	0.10
Urinary Tract Infections	10.67	10.70	11.01	10.60	10.01
Weight Loss	15.45	15.65	15.14	15.80	13.83
Dehydration	0.84	0.84	0.82	0.83	0.93
Bedfast Residents	4.81	5.01	5.04	4.54	3.20
Little/No Activity	15.67	15.86	15.82	15.34	14.32
Stage 1-4 Pressure Ulcer	15.48	16.69	14.11	13.52	11.24

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.



# Region VI – AR, LA, NM, OK, TX



Resident Characteristics*	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
Nursing Home Residents	167,985	88,441	32,396	31,874	15,017
Percent of Nursing Home Residents	100.00	52.65	19.29	18.97	8.94
Percent Female	72.36	72.46	72.37	71.81	72.86
Age: <25	0.26	0.35	0.13	0.13	0.22
Age: 25-54	3.91	4.67	3.30	2.84	3.13
Age: 55-64	5.31	5.74	4.96	4.83	4.53
Age: 65-74	12.43	13.22	12.10	11.68	10.17
Age: 75-84	31.44	31.95	31.30	30.80	30.10
Age: 85-94	38.24	36.20	39.40	40.66	42.47
Age: 95+	8.42	7.86	8.81	9.06	9.38
American Indian/Alaskan Native	0.91	0.47	1.76	1.12	1.26
Asian/Pacific Islander	0.27	0.42	0.15	0.03	0.06
Black	13.84	15.68	13.02	12.58	7.64
Hispanic	7.55	9.98	6.11	4.96	1.92
White	77.44	73.46	78.97	81.30	89.12
Lived Alone	26.05	21.39	28.21	32.07	33.57
Lived with Others	59.35	63.31	57.68	54.73	51.57
Lived in Other Facility	14.60	15.29	14.11	13.20	14.86
Admitted from Private Home with No Services	25.14	22.65	24.96	28.56	31.56
Admitted from Private Home with Services	6.33	5.37	6.59	7.37	8.75
Admitted from Assisted Living	3.42	3.87	3.39	2.50	3.03
Admitted from Other Nursing Home	17.35	18.55	15.65	16.20	17.03
Admitted from Acute Care Hospital	40.05	41.43	41.75	38.40	32.46
Admitted from Other Setting	7.71	8.13	7.66	6.97	7.17
CPS: Intact or Borderline Intact	24.43	23.58	25.06	26.17	24.51
CPS: Mild to Moderately Severely Impaired	48.82	49.56	49.25	47.06	47.00
CPS: Severely or Very Severely Impaired	26.75	26.85	25.69	26.77	28.49
ADL: Independent or Needs Supervision	16.25	15.51	15.86	17.77	18.27
ADL: Limited Physical Assistance	14.78	13.81	16.39	15.41	15.66
ADL: Extensive Assistance in Early Loss ADLs	19.91	20.94	18.98	19.04	17.58
ADL: Extensive Assistance in Later Loss ADLs	9.30	8.91	9.95	9.49	9.78
ADL: Totally Dependent in Some ADLs	19.00	19.67	18.82	17.78	18.00
ADL: Totally Dependent in All Four ADLs	20.77	21.17	20.00	20.51	20.71



## Region VI 2001 (continued)



	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
CHES: No Instability	40.40	42.69	37.55	38.27	37.63
CHES: Slightly Unstable	28.71	28.26	28.95	29.61	28.96
CHES: Mildly Unstable	17.69	16.79	19.27	18.12	18.69
CHES: Moderately Unstable	8.80	8.16	9.72	9.47	9.24
CHES: Highly Unstable	4.39	4.11	4.52	4.54	5.48
Total Case-Mix Index (average)	0.87	0.88	0.87	0.85	0.83
Nursing Case-Mix Index (average)	0.79	0.79	0.79	0.78	0.78
<b>Quality Indicators</b>					
Depression without Therapy	5.60	4.82	6.19	6.58	6.86
9+ Different Medications	47.82	47.91	46.58	48.11	49.25
Bladder/Bowel Incontinence without Plan	66.38	67.31	65.09	65.58	66.78
Indwelling Catheters	11.07	11.21	11.03	11.15	10.14
Tube Feeding	9.79	11.17	8.61	8.17	7.63
Antipsychotic Use without Conditions	22.12	23.03	21.29	20.92	21.18
Antianxiety/Hypnotic Use	20.48	20.46	19.96	20.34	22.02
Daily Physical Restraint	18.94	18.90	18.09	19.78	19.33
Falls	14.48	14.09	15.39	14.48	14.80
Behavioral Symptoms Affecting Others	19.24	18.43	19.77	20.43	20.34
Symptoms of Depression	10.98	9.84	11.76	12.29	13.24
Bladder/Bowel Incontinence	58.05	60.48	56.47	54.68	54.26
Fecal Impaction	0.47	0.39	0.44	0.57	0.72
Urinary Tract Infections	10.30	10.38	9.97	10.12	10.89
Weight Loss	17.36	17.18	18.04	17.27	17.18
Dehydration	1.30	1.19	1.25	1.59	1.43
Bedfast Residents	10.89	10.89	10.14	11.79	10.56
Little/No Activity	24.84	23.64	24.14	26.52	29.87
Stage 1-4 Pressure Ulcer	17.08	17.67	17.03	16.39	15.01

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.



## Region VII – IA, KS, MO, NE



Resident Characteristics*	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
Nursing Home Residents	119,918	46,297	20,558	24,894	28,055
Percent of Nursing Home Residents	100.00	38.61	17.14	20.76	23.40
Percent Female	73.43	74.38	72.47	73.50	72.44
Age: <25	0.16	0.29	0.13	0.08	0.05
Age: 25-54	3.04	3.95	2.82	2.50	2.20
Age: 55-64	3.54	4.21	3.74	2.83	2.94
Age: 65-74	9.46	10.61	9.63	8.21	8.55
Age: 75-84	28.79	30.43	29.11	28.11	26.49
Age: 85-94	43.48	40.19	43.17	46.04	46.86
Age: 95+	11.52	10.32	11.41	12.22	12.92
American Indian/Alaskan Native	0.29	0.31	0.20	0.32	0.31
Asian/Pacific Islander	0.16	0.26	0.13	0.09	0.06
Black	4.37	9.45	2.32	1.00	0.49
Hispanic	0.53	0.74	0.79	0.24	0.25
White	94.65	89.23	96.55	98.36	98.89
Lived Alone	34.96	29.50	35.08	39.84	40.14
Lived with Others	47.13	51.33	44.60	43.25	44.99
Lived in Other Facility	17.90	19.17	20.32	16.91	14.86
Admitted from Private Home with No Services	21.58	17.80	22.06	24.19	25.54
Admitted from Private Home with Services	8.02	6.29	8.14	8.77	10.22
Admitted from Assisted Living	6.96	7.84	7.36	6.71	5.40
Admitted from Other Nursing Home	15.72	17.33	17.25	13.71	13.55
Admitted from Acute Care Hospital	41.28	44.77	39.03	39.20	38.67
Admitted from Other Setting	6.43	5.97	6.16	7.41	6.62
CPS: Intact or Borderline Intact	23.02	21.91	22.34	24.19	24.37
CPS: Mild to Moderately Severely Impaired	54.72	55.54	54.39	53.64	54.50
CPS: Severely or Very Severely Impaired	22.26	22.55	23.27	22.17	21.13
ADL: Independent or Needs Supervision	17.36	16.23	16.78	17.82	19.27
ADL: Limited Physical Assistance	16.86	17.10	17.15	16.73	16.36
ADL: Extensive Assistance in Early Loss ADLs	20.68	21.34	20.38	20.32	20.15
ADL: Extensive Assistance in Later Loss ADLs	13.94	12.60	13.96	15.33	14.92
ADL: Totally Dependent in Some ADLs	19.21	19.79	20.12	18.55	18.13
ADL: Totally Dependent in All Four ADLs	11.95	12.93	11.62	11.24	11.18



## Region VII 2001 (continued)



	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
CHES: No Instability	27.25	29.20	25.71	25.13	27.10
CHES: Slightly Unstable	31.07	30.32	31.75	31.79	31.13
CHES: Mildly Unstable	22.50	21.99	23.30	23.29	22.07
CHES: Moderately Unstable	12.26	11.93	12.46	12.58	12.34
CHES: Highly Unstable	6.92	6.56	6.79	7.21	7.37
Total Case-Mix Index (average)	0.85	0.88	0.86	0.83	0.83
Nursing Case-Mix Index (average)	0.78	0.79	0.79	0.78	0.78
<b>Quality Indicators</b>					
Depression without Therapy	8.89	7.60	9.09	9.43	10.35
9+ Different Medications	49.00	46.61	51.36	49.56	50.74
Bladder/Bowel Incontinence without Plan	24.31	24.95	23.94	24.84	23.18
Indwelling Catheters	8.28	8.96	8.42	7.62	7.64
Tube Feeding	3.69	5.55	3.10	2.41	2.18
Antipsychotic Use without Conditions	17.09	18.35	17.41	16.66	15.23
Antianxiety/Hypnotic Use	16.89	16.45	17.41	17.28	16.86
Daily Physical Restraint	5.92	6.59	4.78	5.51	6.02
Falls	20.30	20.36	21.04	20.35	19.62
Behavioral Symptoms Affecting Others	23.82	22.38	24.57	24.98	24.63
Symptoms of Depression	19.15	17.20	19.16	20.06	21.54
Bladder/Bowel Incontinence	52.95	54.96	51.42	52.63	51.04
Fecal Impaction	0.21	0.21	0.22	0.23	0.17
Urinary Tract Infections	10.64	10.90	10.64	10.52	10.31
Weight Loss	16.01	16.99	16.80	14.91	14.80
Dehydration	0.88	0.81	0.98	0.91	0.88
Bedfast Residents	3.43	4.19	3.17	2.98	2.78
Little/No Activity	18.61	18.84	18.69	19.14	17.58
Stage 1-4 Pressure Ulcer	13.33	16.16	13.73	11.81	10.18

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.



# Region VIII – CO, MT, ND, SD, UT, WY



Resident Characteristics*	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
Nursing Home Residents	51,022	25,314	7,710	7,630	10,368
Percent of Nursing Home Residents	100.00	49.61	15.11	14.95	20.32
Percent Female	70.30	71.18	71.15	67.78	69.37
Age: <25	0.16	0.24	0.06	0.08	0.10
Age: 25-54	3.70	4.96	2.48	2.99	2.07
Age: 55-64	4.04	5.05	3.14	3.20	2.86
Age: 65-74	9.64	10.46	8.85	9.02	8.68
Age: 75-84	30.00	30.67	29.49	30.72	28.22
Age: 85-94	41.95	39.35	44.80	42.99	45.39
Age: 95+	10.51	9.27	11.18	11.01	12.67
American Indian/Alaskan Native	1.76	0.68	0.97	3.34	3.83
Asian/Pacific Islander	0.36	0.61	0.16	0.18	0.06
Black	1.33	2.55	0.16	0.21	0.06
Hispanic	3.15	4.55	0.87	4.07	0.75
White	93.40	91.62	97.85	92.20	95.30
Lived Alone	27.57	23.23	35.22	29.22	32.77
Lived with Others	52.20	56.33	47.13	51.12	43.38
Lived in Other Facility	20.23	20.44	17.66	19.65	23.85
Admitted from Private Home with No Services	16.91	15.47	15.86	20.94	20.74
Admitted from Private Home with Services	5.70	4.71	7.05	7.17	6.44
Admitted from Assisted Living	9.00	9.56	7.23	9.93	8.42
Admitted from Other Nursing Home	15.56	17.67	12.47	13.48	13.23
Admitted from Acute Care Hospital	43.97	44.67	47.66	43.26	36.02
Admitted from Other Setting	8.86	7.92	9.74	5.22	15.15
CPS: Intact or Borderline Intact	22.29	19.74	25.46	23.01	25.63
CPS: Mild to Moderately Severely Impaired	57.67	59.28	55.95	56.38	55.98
CPS: Severely or Very Severely Impaired	20.04	20.98	18.59	20.60	18.39
ADL: Independent or Needs Supervision	16.53	15.88	16.07	16.92	18.17
ADL: Limited Physical Assistance	15.21	16.26	14.12	14.67	13.87
ADL: Extensive Assistance in Early Loss ADLs	22.73	23.34	22.28	21.90	22.19
ADL: Extensive Assistance in Later Loss ADLs	15.01	13.46	17.37	17.00	15.58
ADL: Totally Dependent in Some ADLs	20.04	20.37	20.53	20.24	18.71
ADL: Totally Dependent in All Four ADLs	10.48	10.69	9.62	9.28	11.48



## Region VIII 2001 (continued)



	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
CHES: No Instability	25.61	25.84	24.29	26.12	25.67
CHES: Slightly Unstable	31.74	31.63	31.76	31.13	32.44
CHES: Mildly Unstable	23.65	23.21	24.93	23.59	23.81
CHES: Moderately Unstable	12.64	12.65	12.92	13.29	11.93
CHES: Highly Unstable	6.36	6.67	6.10	5.87	6.15
Total Case-Mix Index (average)	0.87	0.88	0.89	0.84	0.85
Nursing Case-Mix Index (average)	0.80	0.80	0.82	0.80	0.81
<b>Quality Indicators</b>					
Depression without Therapy	10.21	9.06	9.96	12.40	11.58
9+ Different Medications	49.15	46.87	51.35	48.82	53.31
Bladder/Bowel Incontinence without Plan	30.08	29.75	28.95	29.16	32.23
Indwelling Catheters	8.67	9.14	9.56	8.11	7.27
Tube Feeding	3.05	3.90	2.54	2.07	2.10
Antipsychotic Use without Conditions	16.24	17.53	15.50	15.82	13.94
Antianxiety/Hypnotic Use	15.52	15.38	16.73	16.12	14.51
Daily Physical Restraint	8.13	8.53	7.86	7.89	7.51
Falls	23.08	22.76	24.25	23.35	22.77
Behavioral Symptoms Affecting Others	25.21	25.17	23.27	27.25	25.26
Symptoms of Depression	21.63	20.65	20.11	24.58	22.98
Bladder/Bowel Incontinence	54.39	56.49	51.43	53.15	52.39
Fecal Impaction	0.16	0.13	0.14	0.20	0.21
Urinary Tract Infections	10.48	11.07	9.44	9.27	10.68
Weight Loss	15.87	16.75	16.00	15.11	14.19
Dehydration	1.11	0.98	1.06	1.15	1.45
Bedfast Residents	3.53	3.64	3.67	3.42	3.22
Little/No Activity	17.27	18.05	15.19	16.50	17.48
Stage 1-4 Pressure Ulcer	11.68	12.54	11.58	11.66	9.78

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.



# Region IX – AZ, CA, HI, NV



Resident Characteristics*	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
Nursing Home Residents	115,637	105,388	6,405	2,704	815
Percent of Nursing Home Residents	100.00	91.14	5.54	2.34	0.70
Percent Female	68.83	68.86	68.06	68.33	72.64
Age: <25	0.54	0.58	0.22	0.15	0.25
Age: 25-54	7.08	7.33	4.67	4.88	1.72
Age: 55-64	5.67	5.77	4.54	4.77	4.42
Age: 65-74	11.79	11.89	10.69	11.43	8.96
Age: 75-84	30.62	30.46	32.51	30.92	34.72
Age: 85-94	35.42	35.14	38.14	38.79	37.30
Age: 95+	8.88	8.83	9.23	9.06	12.64
American Indian/Alaskan Native	0.83	0.57	1.37	9.61	1.96
Asian/Pacific Islander	8.27	7.98	13.73	3.28	20.12
Black	8.32	8.95	1.83	1.94	0.61
Hispanic	9.77	9.91	10.50	5.22	2.09
White	72.80	72.59	72.57	79.96	75.21
Lived Alone	16.57	16.67	14.65	20.05	17.11
Lived with Others	66.46	66.02	71.67	66.43	66.31
Lived in Other Facility	16.97	17.31	13.69	13.53	16.58
Admitted from Private Home with No Services	11.00	10.57	14.52	15.46	27.27
Admitted from Private Home with Services	3.20	3.03	4.75	6.28	4.28
Admitted from Assisted Living	7.61	7.68	7.41	5.31	6.42
Admitted from Other Nursing Home	13.12	13.36	9.59	20.29	6.95
Admitted from Acute Care Hospital	57.55	57.50	59.33	49.52	50.80
Admitted from Other Setting	7.52	7.85	4.40	3.14	4.28
CPS: Intact or Borderline Intact	20.31	20.22	21.01	21.82	20.12
CPS: Mild to Moderately Severely Impaired	53.21	53.24	53.24	51.63	55.21
CPS: Severely or Very Severely Impaired	26.48	26.54	25.75	26.55	24.66
ADL: Independent or Needs Supervision	10.55	10.48	11.48	10.58	13.87
ADL: Limited Physical Assistance	13.04	13.03	13.01	12.57	15.71
ADL: Extensive Assistance in Early Loss ADLs	21.56	21.47	21.95	23.63	20.98
ADL: Extensive Assistance in Later Loss ADLs	10.45	10.41	10.20	12.87	9.82
ADL: Totally Dependent in Some ADLs	21.83	21.74	23.26	22.12	21.72
ADL: Totally Dependent in All Four ADLs	22.57	22.86	20.11	18.23	17.91



## Region IX 2001 (continued)



	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
CHES: No Instability	41.77	42.33	35.94	36.61	33.50
CHES: Slightly Unstable	29.07	28.93	31.05	28.88	31.90
CHES: Mildly Unstable	16.24	16.10	17.64	18.27	17.67
CHES: Moderately Unstable	8.77	8.64	10.05	9.91	11.04
CHES: Highly Unstable	4.14	4.01	5.31	6.32	5.89
Total Case-Mix Index (average)	0.91	0.91	0.94	0.84	0.82
Nursing Case-Mix Index (average)	0.82	0.82	0.83	0.80	0.77
<b>Quality Indicators</b>					
Depression without Therapy	5.56	5.26	8.70	8.68	9.83
9+ Different Medications	40.61	40.60	40.73	41.09	36.20
Bladder/Bowel Incontinence without Plan	38.45	38.42	33.98	47.28	38.68
Indwelling Catheters	9.55	9.57	9.21	10.39	5.40
Tube Feeding	12.29	12.83	7.56	5.07	3.31
Antipsychotic Use without Conditions	15.20	15.29	14.42	14.70	14.93
Antianxiety/Hypnotic Use	14.22	14.01	16.51	17.04	11.33
Daily Physical Restraint	19.05	19.31	16.13	17.12	13.87
Falls	11.39	11.00	15.75	14.83	16.44
Behavioral Symptoms Affecting Others	19.43	19.31	19.76	22.62	24.57
Symptoms of Depression	10.90	10.36	15.96	18.11	19.53
Bladder/Bowel Incontinence	68.09	68.33	65.48	64.28	66.75
Fecal Impaction	0.13	0.12	0.23	0.22	0.12
Urinary Tract Infections	10.09	10.02	10.98	10.95	8.22
Weight Loss	14.90	14.84	15.19	16.31	15.83
Dehydration	0.94	0.93	1.08	0.96	1.35
Bedfast Residents	8.33	8.24	8.84	10.91	6.13
Little/No Activity	12.69	12.49	13.29	18.55	16.34
Stage 1-4 Pressure Ulcer	18.56	18.80	16.63	16.79	8.82

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.



# Region X – AK, ID, OR, WA



Resident Characteristics*	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
Nursing Home Residents	37,648	26,327	6,917	3,093	1,311
Percent of Nursing Home Residents	100.00	69.93	18.37	8.22	3.48
Percent Female	68.86	68.93	68.44	68.87	69.79
Age: <25	0.39	0.48	0.17	0.32	0.00
Age: 25-54	5.05	5.54	3.21	5.69	3.51
Age: 55-64	4.73	4.86	4.25	4.62	4.73
Age: 65-74	10.79	11.20	10.26	8.79	10.14
Age: 75-84	32.50	32.24	34.64	32.23	27.00
Age: 85-94	37.91	37.23	38.59	39.70	43.94
Age: 95+	8.62	8.45	8.88	8.63	10.68
American Indian/Alaskan Native	1.12	0.68	1.24	3.10	4.50
Asian/Pacific Islander	2.01	2.72	0.36	0.45	0.15
Black	1.90	2.54	0.55	0.23	0.00
Hispanic	1.12	1.05	0.94	2.26	0.61
White	93.86	93.00	96.91	93.95	94.74
Lived Alone	17.75	16.73	16.96	27.67	19.45
Lived with Others	58.54	57.74	61.99	57.53	60.87
Lived in Other Facility	23.71	25.53	21.05	14.79	19.68
Admitted from Private Home with No Services	10.89	10.08	11.04	12.60	19.95
Admitted from Private Home with Services	3.71	3.64	3.61	3.56	5.50
Admitted from Assisted Living	12.85	13.40	11.58	9.73	13.76
Admitted from Other Nursing Home	15.70	16.76	13.42	11.51	14.22
Admitted from Acute Care Hospital	51.92	51.05	56.27	58.63	39.45
Admitted from Other Setting	4.92	5.07	4.09	3.97	7.11
CPS: Intact or Borderline Intact	16.09	15.80	16.37	16.72	18.99
CPS: Mild to Moderately Severely Impaired	60.70	60.49	62.50	60.01	57.06
CPS: Severely or Very Severely Impaired	23.22	23.72	21.14	23.28	23.95
ADL: Independent or Needs Supervision	9.62	9.64	9.06	9.54	12.51
ADL: Limited Physical Assistance	12.55	11.98	13.46	14.71	14.04
ADL: Extensive Assistance in Early Loss ADLs	23.93	24.51	23.15	21.27	22.73
ADL: Extensive Assistance in Later Loss ADLs	15.67	15.24	17.33	15.91	14.80
ADL: Totally Dependent in Some ADLs	22.20	22.25	22.05	22.70	20.82
ADL: Totally Dependent in All Four ADLs	16.02	16.37	14.95	15.87	15.10



## Region X 2001 (continued)

	REGION	LOCALE			
		Urban	Large Town	Small Town	Isolated
CHES: No Instability	27.54	28.74	25.75	23.99	21.36
CHES: Slightly Unstable	30.73	30.43	31.23	31.26	32.80
CHES: Mildly Unstable	22.09	21.87	22.06	22.99	24.49
CHES: Moderately Unstable	12.16	11.59	13.72	13.09	13.04
CHES: Highly Unstable	7.49	7.37	7.24	8.66	8.31
Total Case-Mix Index (average)	0.92	0.92	0.91	0.94	0.88
Nursing Case-Mix Index (average)	0.85	0.85	0.84	0.87	0.85
<b>Quality Indicators</b>					
Depression without Therapy	10.17	9.86	10.55	11.14	12.21
9+ Different Medications	52.04	52.40	50.56	51.12	54.77
Bladder/Bowel Incontinence without Plan	40.20	45.08	28.29	32.73	33.22
Indwelling Catheters	10.60	11.07	9.34	9.99	9.31
Tube Feeding	5.44	6.04	3.85	5.11	2.59
Antipsychotic Use without Conditions	16.40	16.31	17.54	14.44	17.02
Antianxiety/Hypnotic Use	14.55	14.54	14.04	15.32	15.58
Daily Physical Restraint	12.08	12.22	12.46	11.87	7.70
Falls	19.45	19.22	20.11	20.40	18.23
Behavioral Symptoms Affecting Others	22.37	21.38	24.51	24.31	26.42
Symptoms of Depression	24.21	23.74	24.26	27.11	26.42
Bladder/Bowel Incontinence	67.02	67.17	66.80	67.45	64.43
Fecal Impaction	0.23	0.23	0.25	0.13	0.31
Urinary Tract Infections	12.95	13.07	12.75	12.77	11.98
Weight Loss	18.37	18.71	17.00	18.89	17.48
Dehydration	2.63	2.98	1.32	2.49	3.05
Bedfast Residents	7.38	7.82	6.52	6.41	5.42
Little/No Activity	16.58	16.17	16.04	21.42	16.28
Stage 1-4 Pressure Ulcer	15.45	15.98	14.43	13.92	13.26

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.



*THIS PAGE IS BLANK*

# *State Results 2001*

*(in alphabetical order)*